

THE BREMERMAN-DIRICHLET PROBLEM FOR UNBOUNDED DOMAINS OF $\mathbb{C}^n$

ALEXANDRU SIMIONIUC, GIUSEPPE TOMASSINI

ABSTRACT. Given an unbounded strongly pseudoconvex domain Ω and a continuous real valued function h defined on $\partial\Omega$, we study the existence of a (maximal) plurisubharmonic function Φ on Ω such that $\Phi|_{\partial\Omega} = h$.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. The Perron-Bremermann function	3
2.2. The Bremermann-Dirichlet problem	5
3. Existence of continuous solutions for unbounded domains	7
4. Maximal solutions for unbounded strongly convex domains	9
4.1. Existence	9
4.2. Continuity	15
4.3. Some examples	17
5. The case of unbounded strongly pseudoconvex domains	19
6. Existence of a pluriharmonic solution	22
7. A generalization for q -plurisubharmonic functions	22
References	25

1. INTRODUCTION

Let $\Omega \subset \mathbb{C}^n$ be a bounded domain and $h : \partial\Omega \rightarrow \mathbb{R}$ a continuous function. The problem of extending h to Ω by a plurisubharmonic function was considered for the first time by Bremermann in [2]. He proved that, if Ω is strongly pseudoconvex, the upper envelope $u_{\Omega,h}$ of the class of plurisubharmonic functions in Ω which are majored by h on $\partial\Omega$, is a plurisubharmonic extension of h which is continuous at $\partial\Omega$. Later on, Walsh ([10]) showed that $u_{\Omega,h}$ is actually continuous and Bedford and Taylor ([1]) proved that $u_{\Omega,h}$ is a solution of the homogeneous complex Monge-Ampère equation.

2000 *Mathematics Subject Classification.* 32U05, 32T15.

Key words and phrases. plurisubharmonic function, strongly pseudoconvex domain, Perron-Bremermann function.

It is worth observing that, if Ω is not strongly pseudoconvex, the boundary value h cannot be arbitrary (cfr. Lemma 2).

A generalization of the problem for q -plurisubharmonic functions and strongly q -pseudoconvex domains was considered by Hunt and Murray ([4]) and by Slodkowski ([8]).

In this paper we deal with the following *Bremermann-Dirichlet problem* consisting of finding a function $u : \overline{\Omega} \rightarrow \mathbb{R}$, such that

$$(\star) \quad \begin{cases} (i) & u \text{ is upper semicontinuous in } \overline{\Omega} \\ (ii) & u \text{ is plurisubharmonic in } \Omega \\ (iii) & u|_{\text{b}\Omega} = h \end{cases}$$

where $\Omega \subset \mathbb{C}^n$ is an unbounded strongly pseudoconvex domain with a C^2 boundary and $h : \text{b}\Omega \rightarrow \mathbb{R}$ a continuous function.

In this situation the problem may not admit non-trivial solutions even for very simple domains. Indeed, in [7], the authors constructed a continuous function h on the boundary $\text{b}\Omega$ of a paraboloid Ω , satisfying $\inf_{\text{b}\Omega} h = -\infty$, such that the only plurisubharmonic function u satisfying $u \leq h$ on $\text{b}\Omega$ is the function $u \equiv -\infty$.

Thus, in $(\star)$ we assume that $h \geq 0$.

In the first part of the paper we prove that the problem $(\star)$ admits a solution u which is continuous in $\overline{\Omega}$ (cfr. Theorem 6).

In the second part of the paper we study the existence of a maximal solution for the Bremermann-Dirichlet problem. We first consider the case of strongly convex domains. Using an appropriate exhaustive sequence of subdomains of Ω and a decreasing sequence of Perron-Bremermann functions defined on bounded domains we prove the following:

- Let $\Omega \subset \mathbb{C}^n$ be an unbounded strongly convex domain and $h : \text{b}\Omega \rightarrow \mathbb{R}$ a bounded continuous function. Then the Bremermann-Dirichlet problem admits a maximal solution, which is continuous on $\overline{\Omega}$.

(cfr. Proposition 19). If $h : \text{b}\Omega \rightarrow [0, +\infty)$ is not bounded, the continuity of the maximal solution, which is always granted on the boundary $\text{b}\Omega$ (cfr. Proposition 15) is hard to prove in general. In Subsection 4.3 we exhibit some examples where we can obtain the continuity of the maximal solution assuming some hypothesis on the boundary of the domain or on the function h .

The general case of a strongly pseudoconvex domain is treated in Section 5. The existence of the maximal solution and its continuity at the boundary are obtained assuming an additional hypothesis for Ω , which was introduced by Lupaccolu ([5]) in studying the extension problem for CR functions in unbounded domains.

The last section of the paper deals with q -plusubharmonic solutions. We prove that if Ω is an unbounded strongly convex domain and h a bounded real valued continuous function defined on $\partial\Omega$, then the generalized Bremermann-Dirichlet problem for q -plusubharmonic functions admits a maximal solution which is continuous on $\overline{\Omega}$ and $(n - q - 1)$ -plurisuperharmonic in Ω .

2. PRELIMINARIES

2.1. The Perron-Bremermann function. A plurisubharmonic function $u : U \rightarrow \mathbb{R} \cup \{-\infty\}$ on an open set $U \subseteq \mathbb{C}^n$ is an upper semicontinuous function such that its restriction to any complex line L is subharmonic, i.e. for every compact subset $K \subset U \cap L$ and for every harmonic function h on $\text{int } K$ continuous on K such that $h|_{\partial K} \geq u|_{\partial K}$ we have $h \geq u|_K$. If $U \subseteq \mathbb{C}^n$ is an open set, we denote by $\text{Psh}(U)$ the set of plurisubharmonic functions in U and by $\text{Psh}^{(s)}(U)$ the subset of functions $u \in \text{Psh}(U)$ which are semicontinuous on $\overline{U}$.

Proposition 1. *Let $U, V \subset \mathbb{C}^n$ be open sets, $V \subset U$, $u \in \text{Psh}^{(s)}(U)$ and $v \in \text{Psh}^{(s)}(V)$ such that $u|_{\partial V} \geq v|_{\partial V}$. The function $\Psi : \overline{U} \rightarrow \mathbb{R} \cup \{-\infty\}$ given by*

$$\Psi(z) = \begin{cases} \max\{u(z), v(z)\}, & z \in \overline{V} \\ u(z), & z \in \overline{U} \setminus \overline{V}, \end{cases}$$

is plurisubharmonic on U .

Proof. Let $m = \max\{u, v\}$. It is clear that Ψ is upper semicontinuous. Let L be a complex line, $K \subset L \cap \overline{U}$ a compact subset and $h : K \rightarrow \mathbb{R}$ a continuous function, harmonic on $\text{int } K$, such that $h|_{\partial K} \geq \Psi|_{\partial K}$. We have to prove that $h \geq \Psi|_K$. Suppose by contradiction that there exists $z^0 \in K$ such that $h(z^0) < \Psi(z^0)$. Since Ψ is plurisubharmonic on V and on $U \setminus \overline{V}$, we must have $K \cap \partial V \neq \emptyset$. If $z^0 \in V$ (respectively $z^0 \in U \setminus \overline{V}$), by the maximum principle applied to the subharmonic function $m|_L$ on $V \cap L$ (respectively $u|_L$ on $U \cap L \setminus \overline{V}$), we deduce in both cases that there exists $z^1 \in K \cap \partial V$ such that

$$u(z^1) = m(z^1) > h(z^1).$$

This contradicts the maximum principle for $u|_L$, since $h|_{\partial K} \geq \Psi|_{\partial K} \geq u|_{\partial K}$. $\square$

A plurisubharmonic function $u : \Omega \rightarrow \mathbb{R} \cup \{-\infty\}$ is called (*locally*) *maximal* if for every open set $G \subset \Omega$ and every function v plurisubharmonic on G such that $\limsup_{z \rightarrow p} v(z) \leq u(p)$ for all $p \in \partial\Omega$ we have $u \leq v$ on G .

Let $\Omega \subseteq \mathbb{C}^n$ be a domain with C^2 boundary. Let $p \in \partial\Omega$ and U be an open neighbourhood of p in $\mathbb{C}^n$. A C^2 function $\varrho : U \rightarrow \mathbb{R}$ satisfying

$$D \cap \Omega = \{z \in U : \varrho(z) < 0\}$$

and $d\varrho \neq 0$ on $\text{b}\Omega \cap U$ is called a *defining function* on U for $\text{b}\Omega$.

Let $L_p(\varrho)$ denote the Levi form

$$\bar{\partial}\partial\varrho(p) = \sum_{j,k=1}^n \frac{\partial^2 \varrho(p)}{\partial z_j \partial \bar{z}_k} dz_i \wedge d\bar{z}_j$$

of ϱ at p . Ω is called strongly pseudoconvex at p if the restriction of $L_p(\varrho)$ to the tangent hyperplane to $\text{b}\Omega$ at p is positively definite. The definition does not depend on the defining function ϱ . A bounded strongly pseudoconvex domain Ω admits a plurisubharmonic globally defining function in a neighbourhood of $\bar{\Omega}$, i.e. can be written in the form

$$\Omega = \{z \in \mathbb{C}^n : \varrho(z) < 0\},$$

where u is strongly plurisubharmonic in a neighborhood of $\bar{\Omega}$ and $d\varrho \neq 0$ on $\text{b}\Omega$.

We state here, the following

Lemma 2. *Let $\Omega \subset \mathbb{C}^n$, $n \geq 2$, be a domain and H a hyperplane such that $H \cap \Omega \neq \emptyset$. Let Ω_1 be a bounded connected component of $\Omega \setminus H$ and Σ and L such that $\text{b}\Omega_1 = \Sigma \cup L$, where $L \neq \emptyset$ is an open subset of H . If $u \in \text{Psh}^{(s)}(\Omega)$, then*

$$\max_{z \in \bar{\Omega}_1} u(z) = \max_{z \in \Sigma} u(z).$$

Proof. Let $z^0 \in \bar{\Omega}_1$. We choose a complex line $L_1 \subset \bar{\Omega}_1$ passing by z^0 such that $L_1 \cap \Omega$ is a domain D of L_1 with $\text{b}D \subset \Sigma$. Considering the constant function $\phi \equiv M$ on $\bar{D}$, where $M = \max_{z \in \Sigma} u(z)$, ϕ is a harmonic function satisfying $u|_{\text{b}D} \leq \phi|_{\text{b}D}$, therefore $u|_D \leq \phi$ and thus $u(z^0) \leq M$. $\square$

For every continuous function $h : \text{b}\Omega \rightarrow \mathbb{R}$, we define

$$\mathcal{P}_{\Omega,h} = \{u \in \text{Psh}^{(s)}(\Omega) : u|_{\text{b}\Omega} \leq h\}$$

and we denote $u_{\Omega,h} : \bar{\Omega} \rightarrow [-\infty, +\infty)$ the upper envelope of $\mathcal{P}_{\Omega,h}$

$$u_{\Omega,h}(z) = \limsup_{\zeta \rightarrow z} \sup_{u \in \mathcal{P}_{\Omega,h}} u(\zeta).$$

$u_{\Omega,h}$ is called the *Perron-Bremermann function* for Ω and h .

Proposition 3. *Let Ω be a bounded domain in $\mathbb{C}^n$. The Perron-Bremermann function has the following properties:*

- (i) $u_{\Omega,h+c} = u_{\Omega,h} + c$, $\forall c \in \mathbb{R}$;
- (ii) if $h_1 \leq h_2$, then $u_{\Omega,h_1} \leq u_{\Omega,h_2}$;
- (iii) if $h = \sup(h_1, h_2)$, then $u_{\Omega,h} \geq \sup(u_{\Omega,h_1}, u_{\Omega,h_2})$ (and in general $u_{\Omega,h} \neq \sup(u_{\Omega,h_1}, u_{\Omega,h_2})$);
- (iv) $\max_{z \in \bar{\Omega}} |u_{\Omega,h_1} - u_{\Omega,h_2}| = \max_{z \in \text{b}\Omega} |h_1 - h_2|$;
- (v) if $h_\nu \rightarrow h$ in $C^0(\text{b}\Omega)$, then $u_{\Omega,h_\nu} \rightarrow u_{\Omega,h}$ in $C^0(\bar{\Omega})$.

Proof. (i), (ii) and (iii) are obvious. To prove (iv), we put

$$M = \max_{z \in \text{b}\Omega} (h_1(z) - h_2(z))$$

and

$$m = \min_{z \in \text{b}\Omega} (h_1(z) - h_2(z)).$$

Using (i), (ii) and (iii) we get:

$$-\max(|m|, |M|) \leq m \leq |u_{\Omega, h_1} - u_{\Omega, h_2}| \leq M \leq \max(|m|, |M|)$$

and from this it follows the conclusion, since

$$\max(|m|, |M|) = \max_{z \in \text{b}\Omega} |h_1 - h_2|.$$

Property (v) follows immediately from (iv). $\square$

2.2. The Bremermann-Dirichlet problem. Given a domain $\Omega \subset \mathbb{C}^n$ and a continuous function $h : \text{b}\Omega \rightarrow \mathbb{R}$, the *Bremermann-Dirichlet problem* consists of finding a function $u : \overline{\Omega} \rightarrow \mathbb{R}$, such that

$$(\star) \begin{cases} (i) & u \text{ is upper semicontinuous in } \overline{\Omega} \\ (ii) & u \text{ is plurisubharmonic in } \Omega \\ (iii) & u|_{\text{b}\Omega} = h \end{cases}$$

Bremermann proved in [2] that if Ω is a strongly pseudoconvex bounded domain, then the Perron-Bremermann function $u_{\Omega, h}$ is an element of $\mathcal{P}_{\Omega, h}$, which is continuous on $\text{b}\Omega$ and $u_{\Omega, h}|_{\text{b}\Omega} = h$. In [10], Walsh proved that $u_{\Omega, h}$ is actually continuous in $\overline{\Omega}$. Thus, for a strongly pseudoconvex bounded domain Ω , $u_{\Omega, h}$ is a continuous maximal solution of the problem $(\star)$.

About regularity we have the following

Proposition 4. *Let $\Omega \subset \mathbb{C}^n$ be a strongly pseudoconvex bounded domain and $h : \text{b}\Omega \rightarrow \mathbb{R}$ a C^2 function. Then the Perron-Bremermann solution $u_{\Omega, h}$ of the problem $(\star)$ is Lipschitz in $\overline{\Omega}$.*

We first prove the

Lemma 5. *Let $\Omega \subset \mathbb{C}^n$ be a strongly pseudoconvex bounded domain and $h : \text{b}\Omega \rightarrow \mathbb{R}$ a continuous function. If the Perron-Bremermann solution $u_{\Omega, h}$ of the problem $(\star)$ is Lipschitz in a δ -neighbourhood Ω_δ of $\text{b}\Omega$ in $\overline{\Omega}$, for some $\delta > 0$, then $u_{\Omega, h}$ is Lipschitz in $\overline{\Omega}$. Moreover, if Ω is convex, then any Lipschitz constant in Ω_δ is also a Lipschitz constant in $\overline{\Omega}$.*

Proof. Set $u = u_{\Omega, h}$ and let $\delta > 0$ such that u is c -Lipschitz in Ω_δ (i.e. Lipschitz with constant c). Let $z^1, z^2 \in \overline{\Omega}$ with $\|z^2 - z^1\| < \delta/3$ and

put $y = z^2 - z^1$. Define the function $\Psi : \overline{\Omega} \rightarrow \mathbb{R}$ by

$$\Psi(z) = \begin{cases} \max\{u(z), u(z+y) - c\|y\|\}, & z \in \overline{\Omega}, \text{dist}(z, \text{b}\Omega) > \frac{\delta}{2} \\ u(z), & z \in \overline{\Omega}, \text{dist}(z, \text{b}\Omega) \leq \frac{\delta}{2}. \end{cases}$$

Since u is c -lipschitz in Ω_δ , for every $z \in \Omega$ with $\text{dist}(z, \text{b}\Omega) = \frac{\delta}{2}$

$$|u(z+y) - u(z)| < c\|y\|,$$

thus, by the Proposition 1, Ψ is plurisubharmonic in Ω and therefore is a solution of the problem $(\star)$. Using the maximality of u we deduce that

$$u(z^2) - u(z^1) < c\|z^2 - z^1\|.$$

Interchanging z^1 and z^2 we obtain

$$|u(z^2) - u(z^1)| < c\|z^2 - z^1\|.$$

Since every two points in $\overline{\Omega}$ can be connected by a path in $\overline{\Omega}$ made by segments of length less then or equal to $\delta/3$, we get the conclusion. If Ω is convex, the path may be choosen a segment . $\square$

Proof of Proposition 4 Let $\tilde{h} : \overline{\Omega} \rightarrow \mathbb{R}$ be a C^2 extension of h and ϕ a C^2 defining function for $\text{b}\Omega$ which is strongly plurisubharmonic. There exists a constant $a > 0$ such that the function $\Phi^- = \tilde{h} + a\phi$ is plurisubharmonic and the function $\Phi^+ = \tilde{h} - a\phi$ is plurisuperharmonic. Then we have $\Phi^-|_{\text{b}\Omega} = \Phi^+|_{\text{b}\Omega} = h$ and $\Phi^- \leq u \leq \Phi^+$. Since Φ^- and Φ^+ are C^2 , they are c -Lipschitz for some constant $k > 0$. Let $\delta > 0$, $z^1, z^2 \in \overline{\Omega}$ such that $\|z^2 - z^1\| = \delta$ and put $y = z^2 - z^1$. Let $z \in \Omega$ such that $\text{dist}(z, \text{b}\Omega) = \delta$ and $\bar{z} \in \text{b}\Omega$ such that $\|z^0 - z\| = \delta$. Then, if $u = u_{\Omega, h}$,

$$u(z+y) \leq \Phi^+(z+y) \leq \Phi^+(z) + k\delta \leq \Phi^+(\bar{z}) + 2k\delta \leq \Phi^-(z) + 3k\delta.$$

In view of Proposition 1, the function

$$\Psi(z) = \begin{cases} \max\{\Phi^-(z), u(z+y) - 3k\delta\}, & z \in \overline{\Omega}, \text{dist}(z, \text{b}\Omega) > \delta \\ \Phi^-(z), & z \in \overline{\Omega}, \text{dist}(z, \text{b}\Omega) \leq \delta, \end{cases}$$

is plurisubharmonic in Ω , hence it is a solution of the problem $(\star)$. Using the maximality of the Bremermann solution u , we get that

$$u(z^2) - u(z^1) < k\|z^2 - z^1\|$$

and interchanging z^1 and z^2 we obtain

$$|u(z^2) - u(z^1)| < c\|z^2 - z^1\|.$$

Thus, $u_{\Omega, h}$ is Lipschitz on Ω_δ , whence the conclusion in view of Lemma 4. $\square$

3. EXISTENCE OF CONTINUOUS SOLUTIONS FOR UNBOUNDED DOMAINS

In this section we want to study the Bremermann-Dirichlet problem $(\star)$, notably the existence of maximal solutions, when Ω is an unbounded domain in $\mathbb{C}^n$. It is worth to observe that the problem $(\star)$ may not admit solutions even for very simple domains. Ideed, in [7] it is proved that if Ω is the strongly convex paraboloid

$$\Omega = \{(z, w) \in \mathbb{C}^2 : \operatorname{Im} w > |z|^2 + (\operatorname{Re} w)^2\},$$

it is possible to construct a continuous function $h : \partial\Omega \rightarrow \mathbb{R}$ such that the only plurisubharmonic function u on Ω which satisfy

$$\limsup_{\zeta \rightarrow z} u(\zeta) \leq h(z), \quad \forall z \in \Omega$$

is the function $u \equiv -\infty$. The reason is that the function h constructed there is negative in a "large part of $\partial\Omega$ ". Thus in the sequel we assume that the boundary value for the problem $(\star)$ is non negative.

We start by proving the following existence theorem

Theorem 6. *Let $\Omega \subset \mathbb{C}^n$ be a strongly pseudoconvex domain and $h : \partial\Omega \rightarrow [0, +\infty)$ a continuous function. Then the problem $(\star)$ admits a solution u which is continuous on $\overline{\Omega}$.*

Proof. For every $z \in \partial\Omega$ there exist a neighbourhood U_z of z and a biholomorphism $\phi_z : U_z \rightarrow U'_z$ such that $\phi_z(U_z \cap \partial\Omega)$ is strongly convex. We take two concentric balls $B'_z \subset C'_z \subset U'_z$ centered at $z' = \phi_z(z)$ and we define a continuous function $f'_z : U'_z \cap \phi(\partial\Omega) \rightarrow [0, 1]$ such that

$$f'_z|_{\phi(\partial\Omega) \cap B'_z} \equiv 1, \quad f'_z|_{U'_z \cap \phi(\partial\Omega) \setminus C'_z} \equiv 0.$$

Put

$$S'_z = \{w \in U'_z \cap \phi(\partial\Omega) : f'_z(w) > 0\}.$$

If $S_z = \phi_{z_j}^{-1}(S'_z)$, then $\{S_z\}_{z \in \partial\Omega}$ is an open covering of $\partial\Omega$. From it, we extract a locally finite covering $\{S_{z_j}\}_{j \in J}$. If $f = f' \circ \phi_{z_j}$, we define the functions $\theta_{z_j}, \theta : \partial\Omega \rightarrow [0, +\infty)$ by

$$\theta_{z_j}(z) = \begin{cases} f(z), & z \in S_{z_j} \\ 0, & \text{otherwise} \end{cases}$$

and $\theta = \sum_{j \in J} \theta_{z_j}$. Since $\{S_{z_j}\}_{j \in J}$ is locally finite, $\theta < +\infty$, thus $\{\theta_{z_j}/\theta\}_{j \in J}$ is a partition of the unity. We define $h_{z_j} : \partial\Omega \rightarrow [0, +\infty)$ by $h_{z_j} = h\theta_{z_j}/\theta$, thus $\sum_{j \in J} h_{z_j} = h$. Putting $h'_{z_j} = h_{z_j} \circ \phi_{z_j}^{-1}$ and $G'_{z_j} = \phi_{z_j}(\overline{\Omega}) \cap C'_{z_j}$, we define $g'_{z_j} : \partial G'_{z_j} \rightarrow [0, +\infty)$ by

$$g'_{z_j}(z') = \begin{cases} h'_{z_j}(z'), & z' \in S'_{z_j} \\ 0, & \text{otherwise.} \end{cases}$$

Notice that G'_{z_j} may not be with C^2 boundary. For all that, we may assume G'_{z_j} to be with C^2 boundary, using an approximation result

proved in [9] (see also [6]), which says that for every neighbourhood of the set of singularities of the boundary of a strongly convex analytic polyhedron there exists a C^2 strongly convex subdomain whose boundary coincides with the boundary of the analytic polyhedron outside of that neighbourhood. We consider the subset

$$D'_{z_j} = \{z' \in \overline{G'_{z_j}} : \exists L_{z'} \ni z' \text{ complex line s.t. } L_{z'} \cap b G'_{z_j} \subseteq b C'_{z_j}\}.$$

By the maximum principle, for every non-negative function $u' \in \text{Psh}^{(s)}(G'_{z_j})$ with $u'|_{b G'_{z_j}} \leq g'_{z_j}$ we have $u'|_{D'_{z_j}} \equiv 0$. Thus, if Γ'_{z_j} is the solution for the Bremermann-Dirichlet problem for the domain G'_{z_j} and the assigned values g'_{z_j} , then $\Gamma'_{z_j}|_{D'_{z_j}} \equiv 0$, thus if $\Gamma_{z_j} = \Gamma'_{z_j} \circ \phi_{z_j}^{-1}$ and $D_{z_j} = \phi_{z_j}^{-1}(D'_{z_j})$, we have $\Gamma_{z_j}|_{D_{z_j}} \equiv 0$. Since by construction D'_{z_j} is a neighbourhood of $b C'_{z_j} \cap \phi_{z_j}(\overline{\Omega})$ in $\overline{G'_{z_j}}$, D_{z_j} is a neighbourhood of $C_{z_j} = \phi_{z_j}^{-1}(C'_{z_j})$ in $\overline{G_{z_j}}$, therefore the function $\psi_{z_j} : \overline{\Omega} \rightarrow \mathbb{R}$ given by

$$\psi_{z_j} = \begin{cases} \Gamma_{z_j}, & \text{on } \overline{\Omega} \cap C_{z_j} \setminus D_{z_j} \\ 0, & \text{otherwise} \end{cases}$$

is plurisubharmonic on Ω . We define on $\overline{\Omega}$ the function u by

$$u = \sum_{j \in J} \psi_{z_j},$$

which is a continuous solution of $(\star)$. $\square$

Remark 7. For every neighbourhood U of $b\Omega$, u may be constructed such that $u|_{\Omega \setminus U} \equiv 0$.

Corollary 8. *If $\Omega \subset \mathbb{C}^n$ is a strongly pseudoconvex domain, then there exists a function $\varrho \in C^0(\overline{\Omega}) \cap \text{Psh}(\Omega)$ such that $\{\varrho = 0\} = b\Omega$.*

Proof. If we apply the previous theorem taking $h \equiv 1$, then the function $\phi : \overline{\Omega} \rightarrow \mathbb{R}$ given by $\phi = u - 1$ satisfy the requirements. Indeed, assume that there exist $z^0 \in \Omega$ such that $u(z^0) \geq 1$. There exist $k \in \mathbb{N}$ and $J_1 = \{j_1, \dots, j_k\} \subset J$ such that $\psi_{z_j}(z^0) = 0$ for every $j \in J \setminus J_1$. Let $W_{z_j} = \text{supp } \psi_{z_j}$ for $j \in J$. By construction, $b W_{z_j} = S_{z_j} \cup T_{z_j}$, such that $\psi_{z_j}|_{T_{z_j}} \equiv 0$ for all $j \in J_1$. On $\bigcup_{j \in J_1} W_{z_j}$ we define $u_1 = \sum_{j \in J_1} \psi_{z_j}$. If we put $W = \text{supp } u_1$, then $b W = S \cup T$, where $S = \bigcup_{j \in J_1} S_{z_j}$ and $T \subset \bigcup_{j \in J_1} T_{z_j}$ such that $\psi_{z_k}|_T \equiv 0$ for all $j \in J_1$. Therefore $u_1|_S \leq h \equiv 1$ and $u_1|_T \equiv 0$ which contradicts the maximum principle for the plurisubharmonic function u_1 . $\square$

"Par abus de langage" also φ will be called a *continuous defining function* for $b\Omega$.

4. MAXIMAL SOLUTIONS FOR UNBOUNDED STRONGLY CONVEX DOMAINS

The first case that we treat is the one of unbounded strongly convex domains of $\mathbb{C}^n$.

4.1. Existence. Let $\Omega \subset \mathbb{C}^n$ be a strongly convex domain, $h : \text{b}\Omega \rightarrow [0, +\infty)$ a continuous function and consider the problem $(\star)$ for the domain Ω and the function h .

We have the following intuitive fact

Proposition 9. *Let Ω be a convex unbounded domain in $\mathbb{C}^n$ with C^2 boundary. Then there exists $v \in \mathbb{C}^n \setminus \{0\}$ such that for every $z \in \Omega$ and for every $k > 0$, we have $z + kv \in \Omega$. Moreover, if Ω does not contain straight lines, there exists $v \in \mathbb{C}^n$ with the previous property and a unique complex hyperplane orthogonal to v which is tangent to $\text{b}\Omega$.*

Proof. Let $z^0 \in \Omega$ and let $S(z^0; \eta)$ be a ball of center z^0 . Suppose by contradiction that for every $z' \in S(z^0; \eta)$, the halfline starting in z^0 and passing by z' intersects $\text{b}\Omega$ in some point z ; we define $d(z') = \|z - z^0\|$. We will prove that $d : S(z^0; \eta) \rightarrow (0, +\infty)$ is a continuous function. Let $u \in S(z^0; \eta)$ and let w be the intersection of the halfline starting in z^0 and passing by u with $\text{b}\Omega$ and let $\varepsilon > 0$. Let H be the tangent hyperplane to $\text{b}\Omega$ in w . Let y be a point on the segment of extremities z^0 and w , such that $\|y - w\| = \varepsilon_1$, with $\varepsilon_1 < \varepsilon/2$ (see figure 1). There exists $\varepsilon_2 < \varepsilon/2$ such that the cylinder $C(z^0y; \varepsilon_2)$ of radius ε_2 around the segment of extremities z^0 and y to be contained in Ω . Thus, the basis $B(y; \varepsilon_2)$ of the cylinder $C(z^0y; \varepsilon_2)$ which contains the point y is included in the ball $B(w; \varepsilon)$. Decreasing, eventually, ε_2 , we may assume that for every point $y' \in B(y; \varepsilon_2)$, the halfline starting in z^0 and passing by y' intersects H inside the ball $B(w; \varepsilon)$. Taking the intersections of all such halflines with $S(z^0; \eta)$ we get a neighborhood U of u on $S(z^0; \eta)$ and taking the intersections of all such halflines with $\text{b}\Omega$, we get a neighborhood W of w in $\text{b}\Omega$ and $W \subset B(w; \varepsilon)$. Thus, for every $u' \in U$, $d(u')$ differs from $d(u)$ less than ε .

Since Ω is unbounded and convex, there exists a sequence (u_ν) of points $u_\nu \in S(z^0; \eta)$, such that $d(u_\nu) \rightarrow +\infty$, which is a contradiction, since d is continuous and $S(z^0; \eta)$ is compact. Therefore, there exists $v \in \mathbb{C}^n \setminus \{0\}$ such that for every $k > 0$, we have $z^0 + kv \in \Omega$.

Let $z \in \Omega$. Suppose by contradiction that the halfline starting from z and parallel with v intersects $\text{b}\Omega$ in some point w . Since Ω is convex, the tangent hyperplane to $\text{b}\Omega$ in w is not parallel with v , thus it will also intersect the halfline starting from z^0 and parallel with v , which contradicts again the convexity of Ω .

Assume now that Ω contains no straight line and let

$$C = \{u \in \mathbb{C}^n \setminus \{0\} : \forall z^0 \in \Omega, \forall k > 0, z^0 + ku \in \Omega\}.$$




FIGURE 1.

Since Ω is convex, if $u, v \in C$, then $\alpha u + \beta v \in C$, for every $\alpha, \beta \geq 0$ and, since Ω contains no straight line, C is included in an open halfspace. Let $v \in C$ such that the normal hyperplane H to v passing by the origin to be the bounding hyperplane of an open halfspace including C . We may assume for simplicity that $v = (1, 0, \dots, 0)$. Let

$$H_x = \{z = (x_1, y_1, \dots, y_n) \in \mathbb{C}^n : x_1 = x\}.$$

The first case we treat is when there exists $x \in \mathbb{R}$ such that $H_x \cap \Omega = \emptyset$. Let $l = \sup\{x \in \mathbb{R} : H_x \cap \Omega = \emptyset\}$. Then $H_l \cap \partial\Omega \neq \emptyset$ and H_l is tangent to $\partial\Omega$.

Now we treat together the subcase when $H_l \cap \partial\Omega = \emptyset$ (i.e. H_l is asymptotic to $\partial\Omega$) and the case when for every $x \in \mathbb{R}$, $H_x \cap \Omega \neq \emptyset$. Let $w^0 = (w_1, \dots, w_n) \in \Omega$. We will prove that there exists $u \in \mathbb{C}^n \setminus \{0\}$ orthogonal to v such that for every $k > 0$, we have $w^0 + ku \in \Omega$. Assume by contradiction that the hyperplane H_{w^0} passing by w^0 and orthogonal to v intersects $\partial\Omega$ in a compact set M . For every $y \in M$, the tangent hyperplane H_y to $\partial\Omega$ in y intersects the line $\{w^0 + kv : k \in \mathbb{R}\}$ in a point $p^y = (p_1, \dots, p_n)$ with $p_1 < w_1$. This implies that the set $\{z \in \Omega : z^1 \leq w_1\}$ is included in the intersection of all halfspaces given by H_y and w^0 , thus is bounded. This is in contradiction with the assumption of the first case, as well as of the second case. Therefore there exists $u \in \mathbb{C}^n \setminus \{0\}$ orthogonal to v such that for every $k > 0$, $w^0 + ku \in \Omega$. We apply the previous proof to show that this property holds for every $z \in \Omega$ and thus $u \in C$. This is a contradiction, since C is included in the open halfspace bounded by H . $\square$

As for the existence of a maximal solution the natural idea is to construct the maximal solution as limit of a sequence of Perron-Bremermann functions defined on bounded domains considering an appropriate exhaustive sequence of subdomains of Ω .

We may assume that the vector given by the Proposition 9 to be $v = (1, 0, \dots, 0)$ and let $k_0 \in \mathbb{R}$ be such that $H_0 = \{z = (x_1, y_1, \dots, x_n, y_n) \in$

$\mathbb{C}^n : x_1 = k_0$ to be the unique hyperplane orthogonal to v and tangent to $\text{b}\Omega$. We may assume that $k_0 = 0$. Let $(c_\nu)_{\nu>0}$ and $(c'_\nu)_{\nu>0}$ be increasing sequences in $\mathbb{R}$ such that

$$\{z \in \Omega : x_1 < c'_\nu\} \subset B(0; c_\nu) \subset \{z \in \Omega : x_1 < c'_{\nu+1}\}, \quad \forall n \in \mathbb{N},$$

(see figure 2). Let

$$\Omega'_\nu = \{z \in \Omega : l(z) < c'_\nu\}$$

and

$$\Omega_\nu = \Omega \cap B(0; c_\nu).$$

Notice that Ω_ν may not be with C^2 boundary. For all that, we may assume Ω_ν to be with C^2 boundary, using the same approximation result as in the proof of the Theorem 6. Let $A' = \text{b}\Omega'_\nu \cap \text{b}\Omega$ and $B'_\nu = \text{b}\Omega'_\nu \cap H_\nu$. By Lemma 2, if $u \in \text{Psh}^{(s)}(\Omega_\nu)$, then

$$\max_{z \in \overline{\Omega'_\nu}} u(z) = \max_{z \in A'} u(z).$$

Let $A_\nu = \text{b}\Omega_\nu \cap \text{b}\Omega$ and $B_\nu = \text{b}\Omega_\nu \cap \text{b}B(0; c_\nu)$. Let

$$\{\phi : \text{b}\Omega_\nu \rightarrow \mathbb{R} : \phi|_{A_\nu} = h|_{A_\nu}\} = \{\phi_{\nu,j}\}_{j \in J}$$

be the set of all continuous extensions of $h|_{A_\nu}$ to $\text{b}\Omega_\nu$ (see figure 2). For every $\nu \in \mathbb{N}$ and for every $j \in J$, let $\Phi_{\nu,j}$ be the solution of the Bremermann-Dirichlet problem of the domain Ω_ν and function $\phi_{\nu,j}$.

Proposition 10. *For every $\nu \in \mathbb{N}$ and for every $z \in \Omega'_\nu$ there exists $a_{z,\nu} \in \mathbb{R}$, such that $\{\Phi_{\nu,j}(z) : j \in J\} = (-\infty, a_{z,\nu})$ or $\{\Phi_{\nu,j}(z) : j \in J\} = (-\infty, a_{z,\nu}]$.*

Proof. For every $j \in J$ there exists a family of functions $\{\phi_{\nu,t}\}_{t \in \mathbb{R}}$, $\phi_{\nu,t} : \text{b}\Omega_\nu \rightarrow \mathbb{R}$, for every $t \in \mathbb{R}$, which contains $\phi_{\nu,j}$ as element, such that for every $z \in B_\nu \setminus A_\nu$ we have:

- (i) $\phi_{\nu,t}(z)$ is continuous strongly increasing with respect to t ;
- (ii) $\lim_{t \rightarrow -\infty} \phi_{\nu,t}(z) = -\infty$;
- (iii) $\lim_{t \rightarrow +\infty} \phi_{\nu,t}(z) = +\infty$.

By Proposition 3, we get: if $t_1 \leq t_2$, then $\phi_{\nu,t_1} \leq \phi_{\nu,t_2}$ and

$$\max_{z \in \overline{\Omega_{\nu,t}}} |\Phi_{\nu,t_1} - \Phi_{\nu,t_2}| = \max_{z \in \text{b}\Omega_{\nu,t}} |\phi_{\nu,t_1} - \phi_{\nu,t_2}|$$

for all $\nu \in \mathbb{N}$ and $t_1, t_2 \in \mathbb{R}$. Since the function $\phi_{\nu,t}(z)$ is continuous with respect to t , for every $z \in \text{b}\Omega_\nu$, the function $\Phi_{\nu,t}(z)$ is continuous with respect to t , for every $z \in \Omega_\nu$, thus its image is an interval. Assume that $\{\Phi_{\nu,t}(z) : t \in \mathbb{R}\}$ is lower bounded in some $z^0 \in \Omega'_\nu$. Choosing $t_k = -k$, with $k \in \mathbb{N}$, we obtain a sequence of decreasing plurisubharmonic functions $(\Phi_{\nu,k})_k$ and passing to the limit for $k \rightarrow \infty$, we obtain a plurisubharmonic function Ψ_ν such that $\Psi_\nu(z^0) > -\infty$. Let $z^1 \in B_\nu \setminus A_\nu$ and $\{l(z) = c\}$ be a complex hyperplane tangent in z^1 to Ω_ν . Then there exists $\varepsilon_0 > 0$, such that $M = \{z \in \Omega_\nu : l(z) =$




FIGURE 2.

$c + \varepsilon, |\varepsilon| < \varepsilon_0\}$ is a neighborhood of z^1 in Ω_ν which is union of domains in complex hyperplanes $M_\varepsilon = \{z \in \Omega_\nu : l(z) = c + \varepsilon\}$ with boundary $bM_\varepsilon \subset (B_\nu \setminus A_\nu)$. Since $\Psi|_{M_\varepsilon}$ is plurisubharmonic and $\Psi|_{bM_\varepsilon} \equiv -\infty$, we get $\Psi|_M \equiv -\infty$, thus $\Psi \equiv -\infty$, contradiction. Therefore, for $z \in \Omega_\nu$ and for a family $\{\phi_{\nu,t}\}_{t \in \mathbb{R}}$, the set $\{\Phi_{\nu,t}(z) : t \in \mathbb{R}\}$ is an interval which is not lower bounded. Thus the set $\{\Phi_{\nu,j}(z) : j \in J\}$ is union of intervals with the same property, which is also an interval not lower bounded.

Using Lemma 2, we get that for every $z \in \Omega'_\nu$, $\{\Phi_{\nu,j}(z) : j \in J\}$ is upper bounded by $\sup_{z \in A'_\nu} h(z)$. $\square$

Proposition 11. *Using the previous notations, if $\nu_1 < \nu_2$, then for every $z \in \Omega'_{\nu_1}$, $a_{z,\nu_1} \geq a_{z,\nu_2}$.*

Proof. Assume by contradiction that there exists $z^0 \in \Omega'_{\nu_1}$ with $a_{z^0,\nu_1} < a_{z^0,\nu_2}$. Then there exists $j_2 \in J$ such that $a_{z^0,\nu_1} < \Phi_{\nu_2,j_2}(z^0)$. Let $j_1 \in J$,

such that

$$\phi_{\nu_1, j_1}(z) > \Phi_{\nu_2, j_2}(z)$$

for all $z \in B_{\nu_1}$. We get the contradiction since Φ_{ν_1, j_1} is the solution of the Bremermann-Dirichlet problem for the function Φ_{ν_1, j_1} , $\phi_{\nu_1, j_1}(z) > \Phi_{\nu_2, j_2}(z)$ on Ω_{ν_1} , but $\Phi_{\nu_1, j_1}(z^0) < \Phi_{\nu_2, j_2}(z^0)$. $\square$

For every $\nu \in \mathbb{N}$, let us define on $\overline{\Omega}'_\nu$

$$\Phi_\nu = \sup_{j \in J} \Phi_{\nu, j}$$

and let

$$\Phi_\nu^*(z) = \limsup_{\zeta \rightarrow z} \Phi_\nu(\zeta)$$

be the upper semicontinuous envelope of Φ_ν , which is a plurisubharmonic function.

Proposition 12. *Using the previous notations, for every $\nu_0 \in \mathbb{N}$, $\{\Phi_\nu^*(z)\}_{\nu \geq \nu_0}$ is a decreasing sequence of functions, plurisubharmonic on Ω'_{ν_0} and for every $\nu \in \mathbb{N}$, the function $\Phi_{\nu+1}^*$ defined as before is continuous on A_ν and satisfies $\Phi_{\nu+1}^*|_{A_\nu} = h|_{A_\nu}$.*

Proof. By the Proposition 11, for every $\nu_0 \in \mathbb{N}$, the sequence $\{\Phi_\nu(z)\}_{\nu \geq \nu_0}$ is decreasing on Ω'_{ν_0} , thus passing to the lim sup for $\zeta \rightarrow z$ in the points $z \in \Omega'_{\nu_0}$, we get that $\{\Phi_\nu^*(z)\}_{\nu \geq \nu_0}$ has the same property.

For every $\nu \in \mathbb{N}$, Φ_ν was constructed as the sup of a family of continuous functions, therefore it is lower semicontinuous, thus for every $z \in A_\nu$

$$(1) \quad \Phi_{\nu+1}^*(z) = \limsup_{\zeta \rightarrow z} \Phi_\nu(\zeta) \geq \liminf_{\zeta \rightarrow z} \Phi_\nu(\zeta) \geq \Phi_\nu(z) = h(z)$$

Let $c : b\Omega'_{\nu+1} \rightarrow \mathbb{R}$ be a continuous function such that

$$c|_{A_\nu} = h|_{A_\nu} = \Phi_{\nu+1}|_{A_\nu}$$

and

$$c|_{b\Omega'_{\nu+1} \setminus A_\nu} \geq \Phi_{\nu+1}|_{b\Omega'_{\nu+1} \setminus A_\nu}.$$

There exists a continuous function $\Gamma : \overline{\Omega}'_{\nu+1} \rightarrow \mathbb{R}$, harmonic on $\Omega'_{\nu+1}$, such that $\Gamma|_{b\Omega'_{\nu+1}} = c$. For every $j \in J$, $\Phi_{\nu+1, j}$ is plurisubharmonic on $\Omega'_{\nu+1}$ and thus, in particular, subharmonic. We have

$$\Phi_{\nu+1, j}|_{b\Omega'_{\nu+1}} \leq \Phi_{\nu+1}|_{b\Omega'_{\nu+1}} \leq c = \Gamma|_{b\Omega'_{\nu+1}},$$

for all $j \in J$, therefore

$$\Phi_{\nu+1, j}|_{\Omega'_{\nu+1}} \leq \Gamma$$

on $\Omega'_{\nu+1}$ for all $j \in J$. Thus

$$\Phi_{\nu+1}|_{\Omega'_{\nu+1}} \leq \Gamma$$

on $\Omega'_{\nu+1}$. Passing to the $\limsup$ for $\zeta \rightarrow z$ in the points $z \in A_\nu$ and using the continuity of Γ , we get

$$(2) \quad \Phi_{\nu+1}^*|_{A_\nu} \leq \Gamma|_{A_\nu} = h|_{A_\nu}.$$

From (1) and (2) it follows $\Phi_{\nu+1}^*|_{A_\nu} = h|_{A_\nu}$. To prove that $\Phi_{\nu+1}^*$ is continuous on A_ν , it suffices to show that it is lower semicontinuous on A_ν , since $\Phi_{\nu+1}^*$ is upper semicontinuous. Indeed,

$$\liminf_{\zeta \rightarrow z} \Phi_{\nu+1}^*(\zeta) \geq \liminf_{\zeta \rightarrow z} \Phi_{\nu+1}(\zeta) \geq \Phi_{\nu+1}(z) = h(z) = \Phi_{\nu+1}^*(z)$$

for all $z \in A_\nu$, since $\Phi_{\nu+1}$ is lower semicontinuous. $\square$

For every $z \in \overline{\Omega}$ there exists $\nu_0 \in \mathbb{N}$, such that $z \in \overline{\Omega}_{\nu_0}$. Let $\Phi : \overline{\Omega} \rightarrow \mathbb{R}$ be defined by

$$(3) \quad \Phi(z) = \inf_{\nu \geq \nu_0} \Phi_\nu^*(z).$$

As a consequence of the way in which it was constructed, the function Φ extends plurisubharmonically the boundary values prescribing function h and it is maximal among the functions with the same property. So we can conclude the section with the following

Theorem 13. *If $\Omega \subset \mathbb{C}^n$ is an unbounded strongly convex domain and $h : \text{b}\Omega \rightarrow [0, +\infty)$ a continuous function, then the Bremermann-Dirichlet problem $(\star)$ admits a (non negative) maximal solution Φ , i.e. if Φ' is another solution of $(\star)$, then $\Phi' \leq \Phi$.*

Proof. We consider the previous construction of the function Φ from (3). Since h is non negative, Φ_ν is non negative for every $\nu \in \mathbb{N}$, thus Φ is non negative. Φ was constructed as the infimum of a family of upper semicontinuous functions, thus it is upper semicontinuous. Since for every $\nu \in \mathbb{N}$, $\Phi_{\nu+1}^*|_{A_\nu} = h|_{A_\nu}$, it is clear that $\Phi|_{\text{b}\Omega} = h$.

Let $\Phi' \in \text{Psh}^{(s)}(\Omega)$ be a function with $\Phi'|_{\text{b}\Omega} \leq h$, let $z \in \overline{\Omega}$ and let $\nu \in \mathbb{N}$ such that $z \in \overline{\Omega}'_\nu$. Let $j \in J$ such that $\Phi'|_{B_\nu} \leq \phi_{\nu,j}|_{B_\nu}$. Then $\Phi'|_{\Omega'_\nu} \leq \Phi_{\nu,j}|_{\Omega'_\nu}$, thus $\Phi'|_{\Omega'_\nu} \leq \Phi_\nu|_{\Omega'_\nu}$ and $\Phi'|_{\Omega'_\nu} \leq \Phi_\nu^*|_{\Omega'_\nu}$. Since the inequality holds for every $\nu \in \mathbb{N}$ with $z \in \overline{\Omega}'_\nu$, we get $\Phi'|_{\Omega'_\nu} \leq \Phi|_{\Omega'_\nu}$. $\square$

Remark 14. The function Φ from the previous theorem is locally maximal.

Indeed, if $G \subset \Omega$ in an open subset and $v \in \text{Psh}^{(s)}(G)$ such that $v|_{\text{b}G} \leq \Phi|_{\text{b}G}$, we may consider the function $\Phi' : \overline{\Omega} \rightarrow \mathbb{R}$ given by

$$\Phi'(z) = \begin{cases} \max\{\Phi(z), v(z)\}, & z \in \overline{G} \\ \Phi(z), & z \in \overline{\Omega} \setminus G \end{cases}$$

By Proposition 1, Φ' is plurisubharmonic, therefore it is a solution of $(\star)$. Using the maximality of Φ , $\Phi' \leq \Phi$, therefore $v \leq \Phi$ on G .

4.2. Continuity.

Proposition 15. *Let $\Omega \subset \mathbb{C}^n$ be an unbounded strongly convex domain and $h : \text{b}\Omega \rightarrow [0, +\infty)$ a continuous function. Then the maximal solution of the Bremermann-Dirichlet problem $(\star)$ is continuous on $\text{b}\Omega$.*

Proof. We consider the previous construction of the function Φ from (3) and we have to prove that it is continuous on $\text{b}\Omega$. Since Φ is upper semicontinuous, it remains to prove its lower semicontinuity. Let $z^0 \in \text{b}\Omega$. By Theorem 6, there exists a solution u of the problem $(\star)$ which is continuous on $\overline{\Omega}$. Let $\Psi : \overline{\Omega} \rightarrow \mathbb{R}$ be given by $\Psi = \max\{\Phi, u\}$. Then Ψ is a solution of the problem $(\star)$, thus by the Proposition 13 it follows that $\Phi \geq \Psi$ on $\overline{\Omega}$, which implies

$$\liminf_{z \rightarrow z^0} \Phi(z) \geq \liminf_{z \rightarrow z^0} \Psi(z) \geq \liminf_{z \rightarrow z^0} \Psi_1(z) = \Psi_1(z^0) = h(z^0) = \Phi(z^0).$$

□

Remark 16. Continuity of h can be relaxed. Indeed, if h is only upper semicontinuous, we can approximate it from above by a decreasing sequence of continuous functions h_ν . Thus we obtain a decreasing sequence of plurisubharmonic functions Φ_ν converging to a limit Φ , which is plurisubharmonic and for each $z \in \text{b}\Omega$ such that $\limsup_{z \rightarrow z^0} h(z) = h(z^0)$ we have $\limsup_{z \rightarrow z^0} \Phi(z) = h(z^0)$.

Proposition 17. *If $\Omega \subset \mathbb{C}^n$ is an unbounded strongly convex domain, $h : \text{b}\Omega \rightarrow [0, +\infty)$ a continuous function, $D \subset \Omega$ is a strongly pseudoconvex subdomain such that the maximal solution Φ of the Bremermann-Dirichlet problem $(\star)$ is continuous on $\text{b}D$, then Φ is continuous on $\overline{D}$.*

Proof. Let u denote the solution of the Bremermann-Dirichlet problem for the domain D and given boundary $\Phi|_{\text{b}D}$. It is clear that $\Phi|_D \leq u$. Define

$$\Psi(z) = \begin{cases} u(z), & z \in \overline{D} \\ \Phi(z), & z \in \overline{\Omega} \setminus \overline{D}, \end{cases}$$

By Proposition 1, the function Φ is plurisubharmonic, thus it is a solution of the problem $(\star)$, therefore, by Theorem 13 we get $\Phi \geq \Psi$ and in particular $\Phi|_{\overline{D}} \geq \Psi|_{\overline{D}} = u$. Thus, $\Phi|_{\overline{D}} = u$, which is a continuous function. □

The continuity of the maximal plurisubharmonic solution Φ on $\overline{\Omega}$ is more involved. It heavily depends on the behaviour of the boundary value h . In the rest of this section we treat some particular case which, in particular, allows us to prove continuity for bounded boundary values.

Let

$$M = \{x_1 : z = (x_1, y_1, \dots, y_n) \in \text{b}\Omega\}$$

and $g : M \rightarrow \mathbb{R}$ be defined by

$$(4) \quad g(x) = \max\{h(z) : x_1 \leq x\};$$

then

$$(5) \quad g(x) = \max\{\Phi(z) : x_1 \leq x\}$$

by Lemma 2.

We have

Lemma 18. *If*

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{x} = 0,$$

then the function Φ defined in (3) is continuous.

Proof. Let $z^0 \in \Omega$, $z^0 = (\xi_1, \zeta_1, \dots, \xi_n, \zeta_n)$ and let $\varepsilon > 0$. We define $R : \mathbb{C}^n \rightarrow \mathbb{R}$ by

$$R(z) = -\frac{\varepsilon}{2\xi_1}x_1 - \frac{\varepsilon}{2}.$$

Since

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{x} - \frac{\varepsilon}{2\xi_1} = -\frac{\varepsilon}{2\xi_1} < 0,$$

we get

$$\lim_{x \rightarrow +\infty} g(x) - \frac{\varepsilon}{2\xi_1}x = -\infty,$$

thus there exists $\nu \in \mathbb{N}$ such that

$$(6) \quad \Phi(z) + R(z) < 0, \quad \forall z = (x_1, y_1, \dots, y_n) \in \Omega \setminus \Omega'_{\nu-2}.$$

Since Φ is continuous on A'_ν , for every $z \in A'_\nu$ there exists $\delta_z > 0$ such that for every $z' \in B(z; 2\delta_z)$,

$$|\Phi(z) - \Phi(z')| < \frac{\varepsilon}{4}.$$

$\bigcup_{x \in A'_\nu} B(z; \delta_z)$ is an open covering of the compact set A'_ν , thus there exists a finite subcovering $\bigcup_{i \in I} B(z_i; \delta_{z_i})$. Let $\delta = \min\{\delta_{z_i} : i \in I\}$. We claim that

$$\Phi(z) \geq \Phi(z^0) - \varepsilon,$$

for all $z \in B(z^0; \delta/3)$, i.e. Φ is lower semicontinuous in z^0 , thus continuous. Let $z' \in B(z^0; \delta/3)$ and $y = z^0 - z'$. Let $\Psi : \overline{\Omega} \rightarrow \mathbb{R}$ be given by

$$\Psi(z) = \begin{cases} \max\{\Phi(z), \Phi(z+y) + R(z)\}, & z \in \overline{\Omega}'_{\nu-1}, \text{ dist}(z, \text{b}\Omega) > \frac{\delta}{3} \\ \Phi(z), & z \in \overline{\Omega}'_{\nu-1}, \text{ dist}(z, \text{b}\Omega) \leq \frac{\delta}{3} \\ \Phi(z), & z \in \overline{\Omega} \setminus \overline{\Omega}'_{\nu-1} \end{cases}$$

We will show that Ψ is plurisubharmonic on Ω . Since Φ is plurisubharmonic on Ω , it remains to prove the plurisubharmonicity of Ψ in a neighborhood of points z with $\text{dist}(z, \text{b}\Omega) = \delta/3$ or $z \in B'_{\nu-1}$. Let $z \in \Omega$ with $\text{dist}(z, \text{b}\Omega) \in (\delta/3, 2\delta/3)$. Then there exists $z'' \in \text{b}\Omega$ with

$\|z - z''\| < 2\delta/3$ and thus $\|(z + y) - z''\| < \delta$. There exists $i \in I$ such that $\|z'' - z_i\| < \delta_{z_i}$, therefore $z, z + y \in B(z_i; 2\delta_{z_i})$, thus

$$|\Phi(z) - \Phi(z_i)| < \varepsilon/4, \quad |\Phi(z + y) - \Phi(z_i)| < \varepsilon/4,$$

thus

$$\Phi(z) > \Phi(z + y) - \varepsilon/2.$$

Therefore, in a neighborhood of points z with $\text{dist}(z, \text{b}\Omega) = \delta/3$, Ψ is equal to Φ . Now, let $z \in B'_{\nu-1}$ with $\text{dist}(z, \text{b}\Omega) > \delta/3$. Then, decreasing eventually δ , $z + y \in \Omega'_\nu \setminus \Omega'_{\nu-2}$ and using the relation (6) and the uniform continuity of the function R on $\overline{\Omega}'_\nu \setminus \Omega'_{\nu-2}$, we get that

$$\Phi(z + y) + R(z) < 0 \leq \Phi(z)$$

in a neighborhood of z . Therefore, in a neighborhood of z , Ψ is equal to Φ .

By the Proposition 13, we get that $\Phi \geq \Psi$. Taking $z = z'$, we get that $\Phi(z') \geq \Psi(z') \geq \Phi(z' + y) - \varepsilon = \Phi(z^0) - \varepsilon$. $\square$

As a immediate consequence, we get the following

Proposition 19. *Let $\Omega \subset \mathbb{C}^n$ be an unbounded strongly convex domain and $h : \text{b}\Omega \rightarrow \mathbb{R}$ a bounded continuous function. Then the maximal solution of the Bremermann-Dirichlet problem $(\star)$ is continuous on $\overline{\Omega}$.*

Remark 20. Under the assumption of the previous proposition, (as well as for other less restrictive hypothesis which guarantee the continuity of the maximal solution, see Subsection 4.3), the maximal solution of the Bremermann-Dirichlet problem $(\star)$ satisfies the distributional definition for the homogeneous complex Monge-Ampère equation introduced by Bedford and Taylor (see [1]).

Some variations on the theme are contained in the next subsection.

4.3. Some examples.

Lemma 21. *Let Ω be a strongly convex domain in coordinates choosen like before. Let $m \in \mathbb{N}$, let*

$$K = \{z = (x_1, y_1, \dots, y_n) \in \mathbb{C}^n : m(2m - 1)y_1^2 - x_1^2 < 0, x_1 > 0\}$$

and assume that there exists $u = (\alpha, \beta, 0, \dots, 0) \in \mathbb{C}^n$ such that $\Omega \subseteq K - u$. Using the previous notations, if

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{x^{2m-1}} = 0,$$

then Φ is continuous.

Proof. The proof is analogue with the proof of the Lemma 18. Let $a > 0$. We put

$$k = \frac{\varepsilon}{2(\xi_1 + \alpha + a)^{2m} - 2m(2m-1)(\zeta^1 + \beta)^2(\xi_1 + \alpha + a)^{2m-2}}$$

and we define $R : \mathbb{C}^n \rightarrow \mathbb{R}$ by

$$R(z) = k(m(2m-1)(y_1 + \beta)^2(x_1 + \alpha + a)^{2m-2} - (x_1 + \alpha + a)^{2m}) - \frac{\varepsilon}{2}.$$

The function R has the following properties:

- (i) $R \in \mathbf{Psh}(\mathbb{C}^n)$;
- (ii) $R(z) < -\varepsilon/2$, $\forall z \in K - u$;
- (iii) $R(z^0) = -\varepsilon$.
- (iv) $R(z) < -k(2ax_1^{2m-1} - a^2x_1^{2m-2})$, $\forall z = (x_1, y_1, \dots, y_n) \in K - u$;

Since

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{x^{2m-1}} - \frac{k(2ax^{2m-1} - a^2x^{2m-2})}{x^{2m-1}} = -2ak < 0,$$

we get

$$\lim_{x \rightarrow +\infty} g(x) - k(2ax^{2m-1} - a^2x^{2m-2}) = -\infty,$$

therefore, using the property (iv) and the definition of g by relation (4), there exists $\nu \in \mathbb{N}$ such that

$$(7) \quad \Phi(z) + R(z) < 0,$$

for all $z \in \Omega \setminus \Omega'_{\nu-2}$. The rest of the proof goes as in Lemma 18. $\square$

Proposition 22. *Let $\Omega \subset \mathbb{C}^n$ be an unbounded strongly convex domain in coordinates choosen like before and $h : \mathbb{b} \Omega \rightarrow [0, +\infty)$ a continuous function. If there exist $a, b, c \in \mathbb{R}$ such that $\Omega \subseteq P$, where*

$$P = \{z \in \mathbb{C}^n : ay_1^2 + by_1 + c < x_1\}$$

and there exists $m_0 \in \mathbb{N}$ such that

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{x^{m_0}} = 0,$$

where g is given by (4), then the maximal solution Φ of the Bremermann-Dirichlet problem $(\star)$ is continuous.

Proof. Let $m \in \mathbb{N}$, such that $2m-1 > m_0$ and let

$$K = \{z = (x_1, y_1, \dots, y_n) \in \mathbb{C}^n : m(2m-1)y_1^2 - x_1^2 < 0, x_1 > 0\}.$$

There exists $u = (\alpha, \beta, 0, \dots, 0) \in \mathbb{C}^n$ such that $P \subset K - u$. By the Lemma 21 it follows that Φ is continuous. $\square$

Proposition 23. *Let $\Omega \subset \mathbb{C}^n$ be an unbounded strongly convex domain in coordinates chosen like before and $h : \text{b}\Omega \rightarrow [0, +\infty)$ a continuous function. Let $a > 0$ and*

$$K = \{z \in \mathbb{C}^n : y_1 \in (-\sqrt{2}/a, \sqrt{2}/a)\}.$$

If $\Omega \subseteq K$ and there exists $\alpha < a$ such that

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{e^{\alpha x}} = 0,$$

where g is given by (4), then the maximal solution Φ of the Bremermann-Dirichlet problem $(\star)$ is continuous.

Proof. We take

$$k = \frac{\varepsilon e^{-\alpha \xi_1}}{2\alpha^2 \zeta_1^2 - 4}$$

and

$$R(z) = k e^{\alpha x} (\alpha^2 y^2 - 2) - \varepsilon/2.$$

The condition (iv) becomes

$$R(z) < 2k \left(\frac{\alpha^2}{a^2} - 1 \right) e^{\alpha x_1}$$

for all $z = (x_1, y_1, \dots, y_n) \in K$ and since

$$\lim_{x \rightarrow +\infty} \frac{g(x)}{e^{\alpha x}} + 2k \left(\frac{\alpha^2}{a^2} - 1 \right) = 2k \left(\frac{\alpha^2}{a^2} - 1 \right) < 0,$$

we get

$$\lim_{x \rightarrow +\infty} g(x) + 2k e^{\alpha x} \left(\frac{\alpha^2}{a^2} - 1 \right) = -\infty$$

From now on the proof is similar to that of Lemma 21. □

5. THE CASE OF UNBOUNDED STRONGLY PSEUDOCONVEX DOMAINS

The construction and the arguments used in the case of strongly convex domains in order to obtain the maximal plurisubharmonic function and to get its continuity on $\text{b}\Omega$, remain true in the case of the strongly pseudoconvex domains except for the "preparation theorem" and the construction of the "exhaustive" sequence of hyperplanes which have compact intersection with $\text{b}\Omega$. The domains Ω'_ν , having the property that any plurisubharmonic function $u \in \text{Psh}^{(s)}(\Omega_\nu)$ reaches its maximum on that part of the boundary of Ω'_ν which is contained in $\text{b}\Omega$ (denoted by A'_ν), were obtained in the strongly convex case as the intersection of Ω with a convenient sequence of parallel halfspaces. In the case of the strongly pseudoconvex domains it may be possible for any sequence of hyperplanes to contain elements which divide Ω in two subsets which are both not relatively compact. Therefore, the domains Ω'_ν will be constructed in a different way, namely as envelopes of plurisubharmonicity.

In this section $\Omega \subset \mathbb{C}^n$ will denote a strongly pseudoconvex domain. Let $(\Sigma_\nu)_{\nu \in \mathbb{N}}$ be a sequence of compact subsets $\Sigma_\nu \subset \text{b}\Omega$, such that $\Sigma_\nu \subset \Sigma_{\nu+1}$ for every $\nu \in \mathbb{N}$ and

$$\bigcup_{\nu \in \mathbb{N}} \Sigma_\nu = \text{b}\Omega.$$

Passing, if necessary, to a subsequence, we may find a sequence of balls $B(0; c_\nu)$ with increasing radius c_ν such that if we denote as in the convex case

$$\Omega_\nu = \Omega \cap B(0; c_\nu),$$

$A_\nu = \text{b}\Omega_\nu \cap \text{b}\Omega$ and $B_\nu = \text{b}\Omega_\nu \cap \text{b}B(0; c_\nu)$, then

$$A_{\nu-1} \subset \Sigma_\nu \subset A_\nu,$$

for every $\nu \geq 1$. We define Ω'_ν as the plurisubharmonic envelope of Σ_ν

$$\Omega'_\nu = \{z \in \overline{\Omega}_\nu : f(z) \leq \max_{\zeta \in \Sigma_\nu} f(\zeta), \forall f \in \text{Psh}^{(s)}(\Omega_\nu)\}$$

Differently from what happens in the case of strongly convex domains, a priori we may now have

$$(8) \quad \bigcup_{\nu \in \mathbb{N}} \Omega'_\nu \subsetneq \Omega.$$

which would not guarantee $\Phi < +\infty$. A sufficient condition in order to have equality in (8) is that for every point $z \in \Omega$ there exists an analytic disc $D_z \ni z$ such that $D_z \cap \text{b}\Omega$ to be compact. A sufficient condition for having this was first pointed out by Lupaccioli [5] in studying the extension problem for CR functions in unbounded domains. Precisely,

(L) if $\overline{\Omega}^\infty$ denotes the closure of $\Omega \subset \mathbb{C}^n \subset \mathbb{CP}^n$ in $\mathbb{CP}^n$, then there exists an algebraic hypersurface

$$Z = \{z \in \mathbb{C}^n : P(z) = 0\}$$

such that $Z \cap \overline{\Omega}^\infty = \emptyset$.

Equivalently

(L') there exists a polynomial $P \in \mathbb{C}[z_1, \dots, z_n]$ such that

$$\Omega \subset \{z \in \mathbb{C}^n : |P(z)|^2 > (1 + |z|^2)^{\deg P}\}.$$

Remark 24. There exist unbounded strongly pseudoconvex domains $\Omega \subset \mathbb{C}^n$ which do not verify the condition (L).

Indeed, if we consider a polynomial $Q \in \mathbb{C}[z_1, \dots, z_n]$ and the domain

$$\Omega = \{z \in \mathbb{C}^n : \|z\|^2 + \log \|Q(z)\|^2 < 1\},$$

then Ω is strongly pseudoconvex since its boundary is a level set of a strongly plurisubharmonic function. If Z_Q is the zero set of Q we have $Z_Q \subset \Omega$. Then for every polynomial $P \in \mathbb{C}[z_1, \dots, z_n]$, the zero set Z_P of P and Z_Q intersects in $\mathbb{CP}^n$, i.e. condition (L) is not satisfied.

Remark 25. It is worthy observing that Ω has no defining function. Indeed, assume that ϱ is strongly plurisubharmonic in a neighbourhood of $\overline{\Omega}$ and such that $\Omega = \{\varrho < 0\}$. Then, $\varrho|_{Z_Q}$ is bounded above, thus extendable on the projective closure $\widehat{Z}_Q$ of Z_Q hence constant: a contradiction.

Theorem 26. *If $\Omega \subset \mathbb{C}^n$ is an unbounded strongly pseudoconvex domain which verifies the condition (L) and $h : \text{b}\Omega \rightarrow [0, +\infty)$ a continuous function, then the Bremermann-Dirichlet problem $(\star)$ admits a (non negative) maximal solution Φ , i.e. if Φ' is an other solution of $(\star)$, then $\Phi' \leq \Phi$. Moreover, the function Φ is continuous on $\text{b}\Omega$.*

Proof. The only thing we have to show in order to conclude the proof (by using the methods of the previous section) is that, up to a holomorphic change of coordinates and a holomorphic embedding $E : \mathbb{C}^n \rightarrow \mathbb{C}^N$, for every $z \in \Omega$, there exists a hyperplane passing by z which intersects $\text{b}\Omega$ in a compact set.

Following [5], we divide the proof in two steps.

Step 1. P linear. We consider $\overline{\Omega} \subset \mathbb{CP}^n = \mathbb{C}^n \cup \mathbb{CP}_\infty^{n-1}$, which is disjoint from $Z = \{P = 0\}$. So we can consider new coordinates of $\mathbb{CP}^n$ in such a way that Z is the $\mathbb{CP}^{n-1}$ at infinity. Now Ω is a relatively compact open set of $(\mathbb{C}^n)' = \mathbb{CP}^n \setminus Z$ and $H_\infty = \mathbb{CP}_\infty^{n-1} \cap (\mathbb{C}^n)'$ is a complex hyperplane containing the topological boundary of $\text{b}\Omega$. For every $z \in \Omega$, the complex hyperplane passing by z and parallel to H_∞ intersects $\text{b}\Omega$ in a compact set M_z . Similarly to the case when Ω was strongly convex, we can define an exhaustive sequence Ω'_ν of subsets of Ω such that every function $u \in \text{Psh}^{(s)}(\Omega'_\nu)$ reaches its maximum on $A'_\nu = \text{b}\Omega'_\nu \cap \text{b}\Omega$. For instance, we can take

$$\Omega'_\nu = \bigcup_{\text{dist}(z, H_\infty) > 1/\nu} M_z.$$

Step 2. P generic. We use the Veronese map V to embed $\mathbb{CP}^n$ in a suitable $\mathbb{CP}^N$ in such a way that $V(Z) = L_0 \cap V(\mathbb{CP}^n)$, where L_0 is a linear subspace. The Veronese map V is defined as follows: let d be the degree of P , and let

$$N = \binom{n+d}{d} - 1.$$

Then V is defined by

$$V(z) = V[z^0 : \dots : z_n] = [\dots : w_I : \dots]_{|I|=d},$$

where $w_I = z^I$. If $P = \sum_{|I|=d} \alpha_I z^I$, then $V(Z) = L_0 \cap V(\mathbb{CP}^n)$, where

$$L_0 = \left\{ \sum_{|I|=d} \alpha_I w_I = 0 \right\}.$$

Again we can change the coordinates such that L_0 is the $\mathbb{CP}^{N-1}$ at infinity. We may now find the complex hyperplane as in Step 1. $\square$

6. EXISTENCE OF A PLURIHARMONIC SOLUTION

Similarly to the plurisubharmonic functions, one can define the plurisuperharmonic functions. Precisely, a plurisuperharmonic function $\Phi : \Omega \rightarrow \mathbb{R} \cup \{+\infty\}$ on a domain $\Omega \subseteq \mathbb{C}^n$ is a lower semicontinuous function such that its restriction to any complex line is superharmonic; equivalently, Φ is plurisuperharmonic if and only if $-\Phi$ is plurisubharmonic.

Obviously, we can obtain for plurisuperharmonic functions results which are similar to those we get in the previous sections for plurisubharmonic functions. Thus, if $\Omega \subset \mathbb{C}^n$ is an unbounded strongly convex domain and $h : \text{b}\Omega \rightarrow \mathbb{R}$ a continuous function, then the Bremermann-Dirichlet problem

$$(\star') \quad \begin{cases} (i) & u \text{ is lower semicontinuous in } \overline{\Omega} \\ (ii) & u \text{ is plurisuperharmonic in } \Omega \\ (iii) & u|_{\text{b}\Omega} = h \end{cases}$$

(which for making sense, we assume h upper bounded), admits a minimal solution χ which is continuous on $\text{b}\Omega$. Moreover, if h is bounded, then χ is continuous on $\overline{\Omega}$.

For the solvability of both $(\star)$ and $(\star')$ problems, we assume that h is bounded. In this case, the respective solutions Φ and χ are continuous on $\overline{\Omega}$, satisfy $\Phi \leq \chi$ and for every harmonic solution η of the classical Dirichlet problem we have $\Phi \leq \eta \leq \chi$.

If there exists $z \in \Omega$ such that $\Phi(z) = \chi(z)$, then for the maximum principle applied to the plurisubharmonic function $\Phi - \chi$, we get $\Phi \equiv \chi$, thus there exists a pluriharmonic solution of the problem.

It is well known that boundary values of pluriharmonic functions must satisfy certain compatibility conditions. In [3] it is constructed a differential operator of the third order $\overline{\partial}_{\text{b}}\omega$ such that: on a strongly pseudoconvex hypersurface, the traces h in the sense of distributions of pluriharmonic functions are solutions of $\overline{\partial}_{\text{b}}\omega$; conversely, if Ω is simply connected and $\text{b}\Omega$ is strongly pseudoconvex, every solution of the operator $\overline{\partial}_{\text{b}}\omega$ can be extended to a pluriharmonic function.

7. A GENERALIZATION FOR q -PLURISUBHARMONIC FUNCTIONS

A C^2 function $u : U \rightarrow \mathbb{R}$ on an open set $U \subseteq \mathbb{C}^n$ is called q -plurisubharmonic, $0 \leq q \leq n-1$, if its Levi form $L(u)$ has at least $(n-q)$ non-negative eigenvalues. A generalization of the problem $(\star)$ for q -plurisubharmonic functions was considered by Hunt and Murray in [4]. The authors gave a new definition for q -plurisubharmonic functions, which is also applicable to less regular functions. Precisely,

a q -plurisubharmonic function $u : U \rightarrow \mathbb{R} \cup \{-\infty\}$ on an open set $U \subseteq \mathbb{C}^n$ is an upper semicontinuous function such that for every $(q+1)$ -dimensional complex plane L intersecting U , for every compact subset $K \subset U \cap L$ and for every superharmonic function v on $\text{int } K$, C^2 up to the boundary such that $v|_{\text{b}K} \geq u|_{\text{b}K}$, we have $v|_K \geq u|_K$. Observe that with this terminology, the 0-plurisubharmonic functions are the plurisubharmonic functions. If $U \subseteq \mathbb{C}^n$ is an open set, we denote by $\text{Psh}_q^{(s)}(U)$ the set of upper semicontinuous functions in $\overline{U}$ and q -plurisubharmonic in U .

Given a domain $\Omega \subset \mathbb{C}^n$ and a continuous function $h : \text{b} \Omega \rightarrow \mathbb{R}$, a generalized Dirichlet problem consists of finding a function $u : \overline{\Omega} \rightarrow \mathbb{R}$, such that

$$(\star'') \quad \begin{cases} (i) & u \text{ is upper semicontinuous in } \overline{\Omega} \\ (ii) & u \text{ is } q\text{-plurisubharmonic in } \Omega \\ (iii) & u|_{\text{b} \Omega} = h \end{cases}$$

Let $\Omega \subseteq \mathbb{C}^n$ be a domain with C^2 boundary. Let ϱ be a defining function for Ω in a neighbourhood U of a point $p \in \text{b} \Omega$. If the Levi form $L_p(\varrho)$ of ϱ at p has at most q negative eigenvalues, Ω is called strongly q -pseudoconvex at p . A bounded strongly q -pseudoconvex domain Ω admits a q -plurisubharmonic globally defining function in a neighbourhood of $\overline{\Omega}$. Hunt and Murray proved that if Ω is a strongly q -pseudoconvex bounded domain and $2q < n$, then the problem $(\star'')$ admits a maximal solution $u_{\Omega, h}^q$; this is the upper envelope of the functions u which are upper semicontinuous on $\overline{\Omega}$, q -plurisubharmonic on Ω and such that $u|_{\text{b} \Omega} \leq h$. Moreover, $u_{\Omega, h}^q$ is continuous on $\overline{\Omega}$ and it is also $(n - q - 1)$ -plurisuperharmonic. The same is true in a slightly more general situation, namely for bounded strongly r -pseudoconvex domains and q -plurisubharmonic functions with $0 \leq q \leq n - r - 1$, as noticed by Slodkowski in [8]. In the same paper, the author call q -Bremermann a continuous function which is both q -plurisubharmonic and $(n - q - 1)$ -plurisuperharmonic and proved that for a continuous function h and for each $0 \leq q \leq n - 1$ there exists at most one function u continuous on $\overline{\Omega}$, q -Bremermann on Ω and such that $u|_{\text{b} \Omega} = h$. In particular, if Ω is a bounded strongly pseudoconvex domain and h a continuous function, then for each $0 \leq q \leq n - 1$ the problem $(\star'')$ has a unique q -Bremermann solution.

In this section we want to extend to unbounded domains the results concerning the q -plurisubharmonic functions and the problem $(\star'')$, as we did in the previous sections for the plurisubharmonic functions and the problem $(\star)$. We assume again that the vector given by the Proposition 9 is $v = (1, 0, \dots, 0)$ and we repeat for the q -plurisubharmonic functions the construction we made in Section 4 for plurisubharmonic functions. The Propositions 1, 3, 10, 11 hold for q -plurisubharmonic functions with similar proofs. Lemma 2, which

was used to prove the upper boundedness of a family of plurisubharmonic functions, holds also for q -plurisubharmonic functions, since the harmonic function $\phi \equiv M$ is in particular superharmonic. The Proposition 12 holds for q -plurisubharmonic functions with a slightly different proof, namely the function Γ must be chosen plurisuperharmonic instead of harmonic, so we will take Γ to be the solution of the Bremermann-Dirichlet problem for plurisuperharmonic functions for the domain $\Omega'_{\nu+1}$ and boundary value c . All this results allow us to state the following

Theorem 27. *If $\Omega \subset \mathbb{C}^n$ is an unbounded strongly convex domain and $h : \partial\Omega \rightarrow [0, +\infty)$ a continuous function, then for every $1 \leq q \leq n - 1$ the generalized Dirichlet problem $(\star'')$ admits a (non negative) maximal solution Φ , i.e. if Φ' is another solution of $(\star'')$, then $\Phi' \leq \Phi$. Moreover, Φ is continuous on $\partial\Omega$.*

Proof. The first part of the proof follows as for plurisubharmonic functions. To prove the continuity of the function Φ on $\partial\Omega$, we observe that the function furnished by the Theorem 6 is in particular q -plurisubharmonic. $\square$

Theorem 28. *Let $\Omega \subset \mathbb{C}^n$ be an unbounded strongly convex domain in coordinates chosen like before and $h : \partial\Omega \rightarrow [0, +\infty)$ a continuous function. If one of the following conditions hold:*

- (i) $\lim_{x \rightarrow +\infty} \frac{g(x)}{x} = 0$ or in particular h is bounded;
- (ii) there exist $a, b, c \in \mathbb{R}$ such that $\Omega \subseteq P$, where

$$P = \{z \in \mathbb{C}^n : ay_1^2 + by_1 + c < x_1\}$$

and there exists $m_0 \in \mathbb{N}$ such that $\lim_{x \rightarrow +\infty} \frac{g(x)}{x^{m_0}} = 0$;

- (iii) there exist $a > 0$ such that $\Omega \subseteq K$, where

$$K = \{z \in \mathbb{C}^n : y_1 \in (-\sqrt{2}/a, \sqrt{2}/a)\}$$

and there exists $\alpha < a$ such that $\lim_{x \rightarrow +\infty} \frac{g(x)}{e^{\alpha x}} = 0$,

where g is given by (4), then the maximal solution Φ of the generalized Dirichlet problem $(\star'')$ is continuous in $\overline{\Omega}$ and $(n - q - 1)$ -plurisuperharmonic in Ω .

Proof. The proof of the continuity is similar with the case of plurisubharmonic functions. In this situation we use a result of Sadullaev (see [8]) which says that the sum between a q -plurisubharmonic function and a r -plurisubharmonic function is $(q + r)$ -plurisubharmonic and in the particular case $r = 0$ the sum between a q -plurisubharmonic function and a plurisubharmonic function is q -plurisubharmonic.

The proof of the fact that the maximal solution Φ is $(n - q - 1)$ -plurisuperharmonic in Ω is similar to the case of bounded domains. Let

L be a $(q+1)$ -dimensional complex plane intersecting Ω , $K \subset \Omega \cap L$ a compact subset and v a subharmonic function on $\text{int } K$, C^2 up to the boundary, such that $v|_K < \Phi|_K$. We may assume that K is a ball and that the complex plane L is given by $z_1 = 0, z_2 = 0, \dots, z_q = 0$, thus $v(z)$ is actually $v(z_{q+1}, \dots, z_n)$. Let $\tilde{v}(z_1, z_2, \dots, z_n)$ be given by

$$\tilde{v}(z_1, z_2, \dots, z_n) = v(z_{q+1}, \dots, z_n) - k(z_1 \bar{z}_1 + \dots + z_q \bar{z}_q),$$

where $k > 0$ is chosen sufficiently large such that $\tilde{v} < \Phi$ on $\text{b } U$, where $U \subset \Omega$ is still to be chosen strongly pseudoconvex domain with C^2 boundary and with closure in Ω whose intersection with L is K . Since the Levi form of $\tilde{v}$ has $(n-q)$ non-negative eigenvalues, $\tilde{v}$ is q -plurisubharmonic on U if for instance U is chosen an ellipsoid which has L as symmetry plane. Since $\tilde{v} < \Phi$ on $\text{b } U$, using the result similar to the Proposition 1 for q -plurisubharmonic functions we get that the function $\Psi : \bar{\Omega} \rightarrow \mathbb{R}$ given by

$$\Psi(z) = \begin{cases} \max\{\Phi, \tilde{v}\}, & \text{on } U \\ \Phi, & \text{on } \Omega \setminus \bar{U}, \end{cases}$$

is a solution of the problem $(\star'')$. Using the maximality of the function Φ , we get that $\tilde{v}|_U < \Phi|_U$ and thus $v|_K = \tilde{v}|_K < \Phi|_K$. $\square$

REFERENCES

- [1] Bedford, Eric and Taylor, B. A., *The Dirichlet problem for a complex Monge-Ampère equation*, Invent. Math., **37** (1976), no. 1, 1-44. MR 0445006 (56 #3351)
- [2] Bremermann, H. J., *On a generalized Dirichlet problem for plurisubharmonic functions and pseudo-convex domains. Characterization of Šilov boundaries*, Trans. Amer. Math. Soc., **91** (1959), 246-276. MR0136766 (25 #227)
- [3] De Bartolomeis, Paolo and Tomassini, Giuseppe, *Traces of pluriharmonic functions*, Comp. Math. **44** (1981), 29-39 (English).
- [4] Hunt, L. R. and Murray, John J., *q -plurisubharmonic functions and a generalized Dirichlet problem*, Michigan Math. J. **25** (1979), no.3, 299-316 (English). MR 512901 (80b:32018)
- [5] Lupacchiolu, Guido, *Valeurs au bord de fonctions holomorphes dans des domaines non bornés de $\mathbf{C}^n$* , C. R. Acad. Sci. Paris Sér. I Math., **304** (1987), no.3, 67-69, (French, with English summary). MR 878828 (88c:32024)
- [6] Gay, Roger and Sebbar, Ahmed, *Division et extension dans l'algèbre $A^\infty(\Omega)$ d'un ouvert pseudo-convexe à bord lisse de $\mathbf{C}^n$* , Math. Z., **189** (1985), no.3, 421-447 (French). MR 783566 (86e:32020)
- [7] Shcherbina, N. and Tomassini, G., *The Dirichlet problem for Levi-flat graphs over unbounded domains*, Internat. Math. Res. Notices, (1999), no.3, 111-151. MR 1672246 (2000a:32018)
- [8] Ślodkowski, Zbigniew, *The Bremermann-Dirichlet problem for q -plurisubharmonic functions*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), **11**, (1984), no. 2, 303-326. MR 764948 (86a:32038)
- [9] Tomassini, Giuseppe, *Sur les algèbres $A^0(\bar{D})$ et $A^\infty(\bar{D})$ d'un domaine pseudo-convexe non borné*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), **10** (1983), no. 2, 243-256 (French). MR 728435 (85c:32029)

- [10] Walsh, J. B., *Continuity of envelopes of plurisubharmonic functions*, J. Math. Mech., **18** (1968/1969), 143-148. MR 0227465 (37 #3049)

SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI, 56126-PISA, ITALY

E-mail address: `a.simioniuc@sns.it`, `g.tomassini@sns.it`